

MANY TOPOLOGICAL ABELIAN GROUPS HAVE DENSE DIVISIBLE SUBGROUPS

R. C. HOOPER

1. Introduction. If G is a complete, metrizable, topological abelian group such that the groups

$$G^m = \{x \in G \mid my = x \text{ for some } y \in G\}$$

are dense in G , then Theorem 2.1 of this paper shows that G has a dense divisible subgroup K . An example appears in §3 of a group G that is not divisible even though G has a dense divisible subgroup. Also some of the possibilities are examined when G is itself divisible. It should be remembered that the set of all divisible elements in an abelian group forms a subgroup.

2. The existence of dense divisible subgroups. Let G be a complete, metrizable, topological abelian group. Define $G^m = \{x \in G \mid my = x \text{ for some } y \in G\}$. Suppose that G^m is dense in G for all $m > 0$.

Let $\{U_n\}_{n=1}^\infty$ be an open basis of the identity of G such that $U_{n+1} + U_{n+1} \subset U_n$. Choose an open neighborhood V_n of the identity of G such that

$$V_n + \cdots + V_n \subset U_n \\ (n! \text{ times})$$

Next a sequence $\{x_i\}_{i=1}^\infty$ will be defined inductively. Given $x \in G$, we suppose $x_i, 1 \leq i \leq n$ are given with $x_1 = x$ and $((j+1)x_{j+1} - x_j) \in V_{j+1}$ whenever x_j, x_{j+1} are defined. Define x_{n+1} so that $((n+1)x_{n+1} - x_n) \in V_{n+1}$. We can choose such an x_{n+1} since G^{n+1} is dense in G . For each n consider the sequence $\{y_{n,m}\}_{m=n+1}^\infty$ where

$$y_{n,m;m>n} = (m)(m-1) \cdots (n+1)x_m.$$

The claim is that $\{y_{n,m}\}_{m=n+1}^\infty$ is a Cauchy sequence. For consider

$$\begin{aligned} y_{n,m+1} - y_{n,m} &= (m+1)(m) \cdots (n+1)x_{m+1} - (m)(m-1) \cdots (n+1)x_m \\ &= (m)(m-1) \cdots (n+1)[(m+1)x_{m+1} - x_m] \in (m)(m-1) \cdots (n+1)V_{m+1} \\ &\subset V_{m+1}. \end{aligned}$$

Thus $\{y_{n,m}\}_{m=n+1}^\infty$ forms a Cauchy sequence since $U_{m+1} + U_{m+1} \subset U_m$.

Let y_n be the element which $\{y_{n,m}\}_m$ converges to. The claim is that $(n+1)y_{n+1} = y_n$. Given U_j , we may choose m so that $y_{n+1,m}$ and

Received by the editors March 4, 1969.

$y_{n,m}$ satisfy $y_{n+1,m} - y_{n+1} \in U_{j+n+2}$ and $y_{n,m} - y_n \in U_{j+1}$. Thus by the definition of $y_{n,m}$ and $y_{n+1,m}$, $(m)(m-1) \cdots (n+2)x_m - y_{n+1} \in U_{j+n+2}$ and $(m)(m-1) \cdots (n+1)x_m - y_n \in U_{j+1}$. Hence

$$\begin{aligned}(n+1)y_{n+1} - y_n &= (n+1)y_{n+1} - (m)(m-1) \cdots (n+1)x_m \\ &\quad + (m)(m-1) \cdots (n+1)x_m - y_n \in U_{j+1} + U_{j+1} \subset U_j.\end{aligned}$$

Since this is true for all j , we have $(n+1)y_{n+1} = y_n$.

As a consequence $n!y_n = y_1$ for all n and y_1 is divisible for all n .

Furthermore,

$$\begin{aligned}y_1 - x &= (y_1 - y_{1,m}) + (y_{1,m} - y_{1,m-1}) + (y_{1,m-1} - y_{1,m-2}) + \cdots \\ &\quad + (y_{1,3} - y_{1,2}) + (y_{1,2} - x) \in U_1 + U_m + U_{m-1} + \cdots \\ &\quad + U_3 + U_2 \subset U_1 + U_1\end{aligned}$$

since for any $j \geq n$, $y_{n,j+1} - y_{n,j} \in U_{j+1}$, $y_{1,2} = 2x_2$, $y_{1,2} - x = 2x_2 - x_1 \in U_2$ and we may choose m so that $y_1 - y_{1,m} \in U_1$.

What this shows since U_1 can be chosen as small a neighborhood as desired and since x was arbitrary, is that the divisible elements are dense in G .

THEOREM 2.1. *A complete, metrizable topological abelian group G has a dense divisible subgroup if and only if the G^m are dense in G for all $m > 0$.*

PROOF. If the G^m are dense, then it was just shown that G has a dense divisible subgroup. On the other hand if G has a dense divisible subgroup H , $G^m \cap H = H$ is dense in G .

The next theorem gives one example of groups which have a dense divisible subgroup.

THEOREM 2.2. *Let G be a topological abelian group. If G has a dense, cyclic subgroup H and G has no proper open subgroups, then the subgroups G^n are dense for all $n > 0$.*

PROOF. Suppose one of the G^n for $n > 0$ is not dense in G . Then if $H_1 = G^n \cap H$, let H_2 be the closure of H_1 in H . Since H is algebraically isomorphic to the integers and H_2 is not the zero subgroup, H_2 is open in H . Hence since H is dense in G , the closure L of H_2 in G is an open subgroup of G . But since we supposed that G^n is not dense in G , L would be a proper open subgroup of G which is a contradiction.

A special class of groups like those of Theorem 2.1 are those which are divisible as well as being complete and metrizable. If H is a divisible group such that the map $\phi_{1/n}: x \rightarrow (1/n)x$ is well defined and a continuous homomorphism for all $x \in H$, then the map $\phi_{1/n}$ extends to a continuous homomorphism of the completion \bar{G} of H . Naturally

such a G must be torsion free. Any topological vector space is one example of such a group.

Suppose on the other hand G is complete, metrizable, separable, and divisible. Then $\phi_n: G \rightarrow G$ is continuous and onto. Thus by the open mapping theorem [1, p. 99], ϕ_n is open.

If G is also torsion free, then the mapping $\phi_{1/n}: x \rightarrow (1/n)x$ is well defined and is a homomorphism. Since ϕ_n is open, $\phi_{1/n} = \phi_n^{-1}$ is continuous. The solenoidal group Σ_a , $a = (1, 2, 3, \dots)$ [2, p. 406 and p. 388] is one example of such a torsion free group G . The a -dic solenoid Σ_a is compact, connected, divisible, and torsion free. But even though the mappings $\phi_{1/n}$ are continuous for all $n > 0$, Σ_a is not a topological vector space.

The circle group is an example of a connected, complete, metrizable divisible group such that the mappings $\phi_{1/n}$ are not well defined for any $n > 0$.

3. The size of the maximal divisible subgroup of G . If G_1 and G_2 are topological groups and $\phi: G_1 \rightarrow G_2$ is a homomorphism, then ϕ is called almost open [1, p. 14 and p. 89] if for every $x \in G$, $y = \phi(x)$, and neighborhood U of x , the closure of $\phi(U)$, $\text{Cl}(\phi(U))$, is a neighborhood of $\phi(y)$. If $\phi: G_1 \rightarrow G_2$ is almost open and H_1 is a dense subgroup of G_1 , then it can be checked that ϕ restricted to H_1 is almost open.

Suppose that we are given an almost open homomorphism $\phi': H_1 \rightarrow H_2$. If H_1 is dense in G_1 , H_2 is dense in G_2 , and $\phi: G_1 \rightarrow G_2$ is an extension of ϕ' to G_1 , then it can be checked that ϕ is almost open.

Even if the G^n are dense in G , the maps $\phi_n: G \rightarrow G$ do not have to be almost open. To see this, first define on the additive integers a topology called the group series topology. Let a set $S = \{(p_i, M_i)\}_{i=1}^\infty$ of pairs of positive integers be given such that $M_i = 2^i$, the greatest common divisor of $\{p_i\}_{i=n}^\infty$ is 1 for all n , and such that $3 \sum_{i=1}^{j-1} M_i < p_j$ for all j . Define

$$U_n = \left\{ x \mid x = \sum_{i=n}^N m_i p_i \text{ for } |m_i| \leq 2^{-n} M_i, \text{ for } m_i \text{ and } N \text{ integers} \right\}.$$

These U_n satisfy the axioms [1, p. 46] for a basis of the neighborhoods of the identity for a topological abelian group, i.e., $U_n = -U_n$, and $U_{n+1} + U_{n+1} \subset U_n$. For if $x, y \in U_{n+1}$, then $x = \sum_{i=1}^N m_i p_i$ and $y = \sum_{i=1}^N m'_i p_i$ for some integers m_i, m'_i, N such that $|m_i| \leq (1/2^{n+1}) M_i$, $|m'_i| \leq (1/2^{n+1}) M_i$. Thus $x+y$ can be written as $x+y = \sum_{i=1}^N (m_i + m'_i) p_i$ with $|m_i + m'_i| \leq (1/2^n) M_i$. This proves that $x+y \in U_n$ and as a result $U_{n+1} + U_{n+1} \subset U_n$.

If $x \in U_n$, then $x = \sum_{i=1}^N m_i p_i$ for some integers N, m_i such that $|m_i| \leq (1/2^n) M_i$. Thus $-x = \sum_{i=1}^N (-m_i) p_i$ and $|-m_i| = |m_i|$

$\leq (1/2^n)M_i$. Whence $-x \in U_n$ and therefore $-U_n = U_n$.

Furthermore if $x \in U_1$, then there is a unique set of integers $\{m_i\}_i$ with $x = \sum_{i=1}^N m_i p_i$ and $|m_i| \leq M_i$. For if not for some m_i, m'_i, N , and j , $x = \sum_{i=1}^N m_i p_i = \sum_{i=1}^N m'_i p_i$ with $|m_i| \leq M_i$, $|m'_i| \leq M_i$ and j the largest index such that $m_j \neq m'_j$. Then $|(m_j - m'_j)p_j| = |\sum_{i=1}^{j-1} (m_i - m'_i)p_i| \leq \sum_{i=1}^{j-1} 2M_i$. But since the p_i were chosen so that $p_j > 3 \sum_{i=1}^{j-1} M_i$, then we have $|(m_j - m'_j)p_j| > 3 \sum_{i=1}^{j-1} M_i$ which is a contradiction. Henceforth if we write $x = \sum_i m_i p_i$, it will be assumed that $|m_i| \leq M_i$. Since the greatest common divisor of the elements of any U_n is 1, then the smallest subgroup of the integers containing any U_n is the integers. Thus if the topological group H is the integers under the topology having a basis of the identity the U_n , H has no proper open subgroups and H^m is dense in H for all $m > 0$.

If $\phi_2: H \rightarrow H$ is defined by $\phi_2: x \rightarrow 2x$, then the claim is that $\text{Cl}(\phi_2(U_3))$ is not a neighborhood of 0. If $\text{Cl}(\phi_2(U_3))$ is a neighborhood of 0, then we can find some U_j with $j \geq 3$ such that $U_j \subset \text{Cl}(\phi_2(U_3))$.

In particular $p_j \in \text{Cl}(\phi_2(U_3))$ and there is a $c \in \phi_2(U_3)$ such that $p_j - c \in U_{j+1}$. If $c = \sum_{i=1}^N n_i p_i$, all n_i are divisible by 2 since $c \in \phi_2(U_3)$. Also

$$p_j - c \in U_j + \phi_2(U_3) \subset U_2 + U_2 \subset U_1$$

implies that we can write $p_j - c = \sum_{i=1}^N q_i p_i$ with the q_i unique. Since 2 divides n_j , $q_j = 1 - n_j$ is not zero. This contradicts the fact that $p_j - c \in U_{j+1}$. Hence $\text{Cl}(\phi_2(U_3))$ is not a neighborhood of 0 and ϕ_2 is not almost open.

Let G be the completion of H , the integers, under this topology and let K be the maximal divisible group of G . K is dense in G by Theorem 2.1. Suppose $K = G$.

Let $\phi'_2: x \rightarrow 2x$ be the extension of ϕ_2 to G . Since G is divisible, ϕ_2 is onto. But since G is complete and ϕ'_2 is continuous and onto, then ϕ'_2 is an open mapping [1, p. 99]. But then since ϕ'_2 is almost open, the restriction ϕ_2 of ϕ'_2 to H is almost open. This is a contradiction. Hence $K \neq G$.

Thus we have an example of a complete, metrizable group G such that even though the G^n are dense, G is not divisible. The author does not know whether K is always uncountable when G is uncountable.

BIBLIOGRAPHY

1. T. Husain, *Introduction to topological groups*, Saunders, Philadelphia, Penn., 1966.
2. E. Hewitt and K. Ross, *Abstract harmonic analysis*, Academic Press, New York, 1963.

UNIVERSITY OF NEVADA